



Soils Map



State: **Indiana**
 County: **Whitley**
 Location: **27-31N-10E**
 Township: **Union**
 Acres: **190.58**
 Date: **12/19/2024**



Soils data provided by USDA and NRCS.

©2024 AgriData, Inc.

Area Symbol: IN183, Soil Area Version: 27

Code	Soil Description	Acres	Percent of field	Non-Irr Class Legend	Non-Irr Class	Irr Class	Alfalfa hay Tons	Corn Bu	Corn silage Tons	Grass legume hay Tons	Grass legume pasture AUM	Oats Bu	Pasture AUM	Soybeans Bu	Winter wheat Bu
MvC2	Morley loam, 6 to 12 percent slopes, eroded	26.31	13.9%		IIle			115	18	4			8	40	52
Se	Sebewa loam, disintegration moraine, 0 to 1 percent slopes	18.84	9.9%		IIw			133		4			9	37	65
RcB	Rawson sandy loam, 2 to 6 percent slopes	18.00	9.4%		Ile			126	18	5			8	44	57
Mg	Mermill loam	15.68	8.2%		IIw			170	23	6			11	49	68
Pw	Pewamo silty clay loam, 0 to 1 percent slopes	13.68	7.2%		IIw			157		5	11			47	64
BmB2	Blount loam, interlobate moraines, 1 to 4 percent slopes, eroded	10.74	5.6%		Ile			137	17	5			9	50	54
BwC	Boyer sandy loam, 6 to 12 percent slopes	9.56	5.0%		IIIle	IIIle	3	90	15	3		60	6	30	47
GsB2	Glynwood loam, 2 to 6 percent slopes, eroded	9.27	4.9%		Ile		5	128	18	4	8	78		41	56

Soils data provided by USDA and NRCS.

Code	Soil Description	Acres	Percent of field	Non-Irr Class Legend	Non-Irr Class	Irr Class	Alfalfa hay Tons	Corn Bu	Corn silage Tons	Grass legume hay Tons	Grass legume pasture AUM	Oats Bu	Pasture AUM	Soybeans Bu	Winter wheat Bu
So	Sloan loam, sandy substratum, 0 to 2 percent slopes, frequently flooded	8.88	4.7%		IIIw			146		5	10			40	
MvB2	Morley loam, 3 to 6 percent slopes, eroded	7.26	3.8%		Ile			119	17	4			8	42	53
ByC3	Boyer loam, 6 to 15 percent slopes, severely eroded	5.88	3.1%		IVe			90		3			6	32	45
KaA	Kalamazoo sandy loam, 0 to 2 percent slopes	5.70	3.0%		IIs			117	16	4			8	41	59
BwB	Boyer sandy loam, 1 to 6 percent slopes	5.28	2.8%		IIIs	Ile	4	98	15	3		62	6	34	51
Co	Coesse silty clay loam	4.84	2.5%		IIw			150	22	5			10	44	60
KaB	Kalamazoo sandy loam, 2 to 6 percent slopes	4.81	2.5%		Ile			117	16	4			8	41	59
Gw	Granby loamy sand	4.37	2.3%		IVw			135	20	5			9	28	54
MxC3	Morley clay loam, 6 to 12 percent slopes, severely eroded	4.09	2.1%		IVe			105	15	4	7			37	47
HbA	Haskins loam, 0 to 3 percent slopes	3.95	2.1%		IIw			158		5	11			59	62
MvD2	Morley loam, 12 to 20 percent slopes, eroded	3.00	1.6%		IVe			100	15	3			7	35	45
Re	Rensselaer loam, 0 to 1 percent slopes	2.29	1.2%		IIw			167		6			11	49	68
RcC	Rawson sandy loam, 6 to 12 percent slopes	2.08	1.1%		IIle			130	20	4			9	46	59
Ms	Milford silty clay loam, 0 to 2 percent slopes	1.79	0.9%		IIw			154		5			11	43	62
MxD3	Morley clay loam, 12 to 18 percent slopes, severely eroded	1.75	0.9%		VIe			92	15	3	7			32	42
Wa	Walkkill silty clay loam	1.55	0.8%		IIIw			165	23	5			11	49	66
Px	Pits, gravel	0.98	0.5%												
Weighted Average					*-	*-	0.5	129	12.7	4.4	2.1	8.5	6.6	41.2	54.1

*- Non Irr Class weighted average cannot be calculated on the current soils data due to missing data.

*- Irr Class weighted average cannot be calculated on the current soils data due to missing data.